

Algebra Prelim

January 5, 2011

- Provide proofs for all statements, citing theorems that may be needed.
- If necessary you may use the results from other parts of this test, even though you may not have successfully proved them.
- Do as many problems as you can and present your solutions as carefully as possible.

Good luck!

- (a) Let V be a 3-dimensional vector space over a field K with basis $\{v_1, v_2, v_3\}$. Find a condition on the characteristic of K that guarantees that $\{v_1 + v_2, v_2 + v_3, v_3 + v_1\}$ is also a basis of V .

(b) If V is a 4-dimensional vector space with basis $\{v_1, v_2, v_3, v_4\}$, show that $\{v_1 + v_2, v_2 + v_3, v_3 + v_4, v_4 + v_1\}$ is not a basis of V .

(c) Conjecture a generalization to the case when $\{v_1 + v_2, v_2 + v_3, \dots, v_n + v_1\}$ is a basis of V . In other words, for which n and which characteristic of K is $\{v_1 + v_2, v_2 + v_3, \dots, v_n + v_1\}$ a basis of V (where $\{v_1, \dots, v_n\}$ is a basis of V)? (You do not need to prove your conjecture.)
- Let $f : V \rightarrow V$ be an endomorphism (linear transformation) of a finite-dimensional vector space V such that $f \circ f = f$, $\ker f \neq \{0\}$, and $\operatorname{im} f \neq \{0\}$.

(a) Find the eigenvalues of f .

(b) Show that f is diagonalizable and describe its diagonal matrix representation as closely as possible.
- Let g be an element of a group G . If there are exactly two conjugates of g in G , show that either there are exactly two conjugates of g^2 in G or g^2 is in the center of G .
- Let G be a cyclic group of order 168 that is generated by a . Consider the group homomorphism $\varphi : G \rightarrow G$, defined by $\varphi(a) = a^{105}$. Determine the order of $\ker \varphi$ and $\operatorname{im} \varphi$.
- Find a generator of the ideal $I = (85, 1 + 13i)$ of $\mathbb{Z}[i] \subset \mathbb{C}$.

6. Let R be a factorial subring (UFD) of \mathbb{C} that does not contain $\sqrt{-5}$. Show that the rings $R[\sqrt{-5}]$ and $R[X]/(X^2 + 5)$ are isomorphic. (Notice that R is not necessarily a field.)
7. Let R be a commutative ring with identity 1, and let I, J be ideals of R such that $I + J = R$. Without using the Chinese Remainder Theorem show:
- (a) $IJ = I \cap J$.
 - (b) There are elements $a \in I$ and $b \in J$ such that $a \equiv 1 \pmod{J}$ and $b \equiv 1 \pmod{I}$.
 - (c) The rings R/IJ and $R/I \oplus R/J$ are isomorphic.
8. Consider $\alpha := \sqrt{2 + \sqrt{3}} \in \mathbb{R}$.
- (a) Determine the minimal polynomial f of α over \mathbb{Q} .
 - (b) Show that f splits over $\mathbb{Q}(\alpha)$.
 - (c) Determine abstractly the Galois group of f over \mathbb{Q} .
9. Let E/K be a Galois extension of degree 55 whose Galois group is not abelian. Determine the number of intermediate fields L with $K \subset L \subset E$ and $[L : K] = 11$.