

## Math 241 - Solutions to Quiz 6- Thursday, December 1

1. Let  $C$  be the semi-ellipse  $x^2 + 4y^2 = 1, x \geq 0$  and consider the vector field

$$F(x, y) = (2, x^2).$$

- (a) Find a parametrization  $\mathbf{r}(t)$  of  $C$  and normal vector  $\mathbf{n}(t)$  pointing to the outside of the ellipse. (3 points)

**SOLUTION:**

$$\mathbf{r}(t) = (\cos t, 1/2 \sin t), \quad -\pi/2 \leq t \leq \pi/2$$

$$\mathbf{r}'(t) = (-\sin t, 1/2 \cos t), \quad -\pi/2 \leq t \leq \pi/2$$

$$\mathbf{n}(t) = (1/2 \cos t, \sin t), \quad -\pi/2 \leq t \leq \pi/2$$

We see that  $\mathbf{n}(t)$  is normal to the curve since  $\mathbf{r}'(t) \cdot \mathbf{n}(t) = 0$  for every  $t$ . It points outward since it is never 0 and  $\mathbf{n}(0) = (1/2, 0)$  points outward.

- (b) Find the flux of  $F$  across  $C$ , moving from inside the ellipse to outside the ellipse. If you couldn't do (a), use  $\mathbf{r}(t) = (\cos t, \sin t)$  and  $\mathbf{n}(t) = (\cos t, \sin t)$ . (3 points)

**SOLUTION:**

Flux of  $F$  across  $C$  from inside to outside =  $\int_C F \cdot \mathbf{u} ds$ , where  $\mathbf{u}$  is an outward pointing unit normal. For the choice of  $\mathbf{n}$  in (a),  $\mathbf{u} = \mathbf{n}/|\mathbf{n}|$  and  $|\mathbf{n}| = |\mathbf{r}'|$  so

$$\begin{aligned} \int_C (2, x^2) \cdot \frac{\mathbf{n}}{|\mathbf{n}|} ds &= \int_0^{2\pi} (2, \cos^2 t) \cdot \frac{(1/2 \cos t, \sin t)}{|\mathbf{r}'(t)|} |\mathbf{r}'(t)| dt \\ &= \int_{-\pi/2}^{\pi/2} \cos t + \cos^2 t \sin t dt = [\sin t - 1/3 \cos^3 t]_{-\pi/2}^{\pi/2} = 1 - (-1) = 2 \end{aligned}$$

2. Let  $F(x, y) = (x^3, 2x^2y)$ .

(a) Find  $\operatorname{div} F$ . (2 points)

**SOLUTION:**

$$\operatorname{div} F = 3x^2 + 2x^2 = 5x^2$$

(b) Use the divergence theorem to compute the flux of  $F$  across the unit circle (with outward pointing normal vector).

**SOLUTION:**

The divergence theorem for curves (another form of Green's Theorem) states

$$\int_C F \cdot \mathbf{n} ds = \int \int_D \operatorname{div} F dA,$$

where  $C$  is the boundary curve of  $D$ . So if  $C$  is the unit circle and  $D$  is the unit disk we have

$$\begin{aligned} \int_C (x^3, 2x^2y) \cdot \mathbf{n} ds &= \int \int_D 5x^2 dA = \int_0^1 \int_0^{2\pi} 5r^2 \cos^2 \theta r dr d\theta \\ &= \left( \int_0^1 5r^3 dr \right) \left( \int_0^{2\pi} \cos^2 \theta d\theta \right) = \frac{5}{4} \pi \end{aligned}$$