## Exam 4

| Name:   | G .:     |
|---------|----------|
| Name:   | Section: |
| 1101110 |          |

Do not remove this answer page — you will return the whole exam. You will be allowed two hours to complete this test. You are allowed to use notes on a single piece of 8.5" x 11" paper, front and back, including formulas and theorems. You are required to turn this page in with your exam. You may use a graphing calculator during the exam, but NO calculator with a Computer Algebra System (CAS). Absolutely no communication device use during the exam is allowed.

The exam consists of 10 multiple choice questions and 5 free response questions. Record your answers to the multiple choice questions on this page by filling in the circle corresponding to the correct answer.

Show <u>all work</u> to receive full credit on the free response problems. It will also help you check your answers to show work on multiple choice problems.

## Multiple Choice Questions

(D)

| 1 | A | $\bigcirc$ B | $\bigcirc$ | D | E |  | 6 | A |
|---|---|--------------|------------|---|---|--|---|---|
|---|---|--------------|------------|---|---|--|---|---|

 $\mathbf{2}$ 

- (A) (B) (C) (D) (E) 7 (A) (B) (C) (D) (E)
- **8** (A) (B) (C) (D) (E) **8** (A) (B) (C) (D) (E)
- **1** (A) (B) (C) (D) (E) **9** (A) (B) (C) (D) (E)
- **5** A B C D E **10** A B C D E

| Multiple |    |    |    |    |    | Total |
|----------|----|----|----|----|----|-------|
| Choice   | 11 | 12 | 13 | 14 | 15 | Score |
| 50       | 10 | 10 | 10 | 10 | 10 | 100   |
|          |    |    |    |    |    |       |
|          |    |    |    |    |    |       |

## Multiple Choice Questions

1. (5 points) If f(0) = 1, f(3) = 2, f'(0) = 8 and f'(3) = 5 and f''(x) is continuous, what is

$$\int_0^3 (x+1)f''(x) \ dx?$$

- A. 11
- B. 60
- C. 18
- D. 80
- E. 16

2. (5 points) What is the best form of the partial fraction decomposition of

$$\frac{13}{(x+1)(x^2-9)}?$$

- A.  $\frac{A}{x+1} + \frac{Bx+C}{x^2-9}$
- B.  $\frac{A}{x+1} + \frac{B}{x-3} + \frac{C}{(x-3)^2}$
- C.  $\frac{A}{x+1} + \frac{B}{x^2 9}$
- D.  $\frac{A}{x+1} + \frac{B}{x+3} + \frac{C}{x-3}$
- E.  $\frac{Ax+B}{x+1} + \frac{Cx+D}{x^2-9}$

3. (5 points) Use the **midpoint rule** with n=4 intervals to approximate  $\int_1^9 \sqrt{1+x^5} \, dx$ .

A. 
$$\frac{2}{3}(\sqrt{1+(1)^5}+4\sqrt{1+(3)^5}+2\sqrt{1+(5)^5}+4\sqrt{1+(7)^5}+\sqrt{1+(9)^5})$$

B. 
$$2(\sqrt{1+(2)^5} + \sqrt{1+(4)^5} + \sqrt{1+(6)^5} + \sqrt{1+(8)^5})$$

C. 
$$\sqrt{1+(1)^5}+2\sqrt{1+(3)^5}+2\sqrt{1+(5)^5}+2\sqrt{1+(7)^5}+\sqrt{1+(9)^5}$$

D. 
$$2(\sqrt{1+(1)^5} + \sqrt{1+(3)^5} + \sqrt{1+(5)^5} + \sqrt{1+(7)^5})$$

E. 
$$2(\sqrt{1+(3)^5} + \sqrt{1+(5)^5} + \sqrt{1+(7)^5} + \sqrt{1+(9)^5})$$

- 4. (5 points) Find the sum of the series  $\sum_{n=1}^{\infty} \frac{2^n + 3^n}{5^n}.$ 
  - A.  $\frac{25}{6}$
  - B. 1
  - C.  $\frac{13}{6}$
  - D.  $\frac{32}{15}$
  - E. This series diverges.

- 5. (5 points) Which of the following series converge?
  - $A. \sum_{n=1}^{\infty} \frac{n^5}{n!}$
  - B.  $\sum_{n=2}^{\infty} \frac{\sqrt{n^6}}{n^4 1}$
  - C.  $\sum_{n=0}^{\infty} \cos(n\pi)$
  - D. all of the given series converge
  - E. none of the given series converge

6. (5 points) What is the radius of convergence for the power series

$$\sum_{n=1}^{\infty} \frac{(-1)^n (x-1)^n}{n^2 3^n} ?$$

- A. 3
- B.  $\infty$
- C. 0
- D. 1
- E.  $\frac{1}{3}$

- 7. (5 points) The line y = 4x + 1 for  $1 \le x \le 3$  is rotated about the x-axis. What is the area of the resulting surface?
  - A.  $72\pi$
  - B.  $\sqrt{17}\pi$
  - C.  $3\sqrt{17}\pi$
  - D.  $18\sqrt{17}\pi$
  - E.  $36\sqrt{17}\pi$

- 8. (5 points) The Cartesian coordinates of a point are (-1,1). Find polar coordinates  $(r,\theta)$  of the point, where r > 0, and  $0 \le \theta < 2\pi$ .
  - A.  $r = \sqrt{2}, \theta = \frac{5\pi}{4}$
  - B.  $r = \frac{1}{\sqrt{2}}, \ \theta = \frac{\pi}{4}$
  - C.  $r = -\sqrt{2}, \ \theta = \frac{3\pi}{4}$
  - D.  $r = \sqrt{2}$ ,  $\theta = \frac{3\pi}{4}$
  - E.  $r = 1, \ \theta = -\frac{\pi}{4}$

9. (5 points) Which of the following integrals computes the **arc length** of the parametric curve  $x(t) = 3 + \cos t$ ,  $y(t) = \ln t$ ,  $4 \le t \le 9$ ?

$$A. \int_4^9 \sqrt{1 - \left(\frac{1}{t \sin t}\right)^2} dt$$

B. 
$$\int_{4}^{9} \sqrt{\sin^2 t + \frac{1}{t^2}} dt$$

C. 
$$\int_{4}^{9} \sqrt{(3+\cos t)^2 + (\ln t)^2} dt$$

D. 
$$\int_{4}^{9} \sin t + \frac{1}{t} dt$$

E. 
$$\int_{4}^{9} 2\pi \ln t \sqrt{1 + \sin^2 t} \ dt$$

10. (5 points) Find the equation of the parabola which has vertex (4,1) and focus (6,1).

A. 
$$x = 8(y-1)^2 + 4$$

B. 
$$y = \frac{1}{8}(x-4)^2 + 1$$

C. 
$$x = \frac{1}{8}(y-1)^2 + 4$$

D. 
$$y = 8(x-4)^2 + 1$$

E. 
$$x = \frac{1}{6}(y-1)^2 - 4$$

Free Response Questions

11. (a) (5 points) Evaluate  $\int \frac{dx}{\sqrt{25-x^2}}$  using trigonometric substitution. Show all steps clearly.

(b) (5 points) Evaluate  $\int \sin^3(x) dx$ . Show all steps clearly.

12. (a) (4 points) Write the Maclaurin series (i.e, the Taylor series centered at x=0) for the function

$$f(x) = \frac{2x^5}{1 - x^6}.$$

(b) (6 points) Does the series  $\sum_{n=6}^{\infty} \frac{(-1)^n}{n-4}$  converge absolutely, converge conditionally, or diverge? Show all steps to justify your answer, and clearly list which test(s) you use.

- 13. Let S be the solid obtained by rotating the region in the first quadrant bounded by  $y = x^3$  and y = 9x about the **y-axis**.
  - (a) (5 points) Set up the integral that computes the volume of S using the  $\mathbf{disk/washer}$  method.

(b) (5 points) Set up the integral that computes the volume of S using the **cylindrical** shells method.

- 14. Consider the polar curve C defined by  $r = \sqrt{6 \sin \theta}$ .
  - (a) (5 points) Set up an integral which computes the area bounded by C which lies in the sector  $0 \le \theta \le \pi$ , and then **evaluate** your integral to find the area.

(b) (5 points) Set up but do **not** evaluate an integral which computes the length of C for  $0 \le \theta \le \pi$ . You do not need to simplify.

15. (a) (5 points) Find the vertices and foci of the hyperbola defined by the equation

$$\frac{(x-3)^2}{1} - \frac{(y+2)^2}{9} = 1$$

(b) (5 points) Find the vertices of the ellipse defined by the equation

$$16x^2 + y^2 - 6y = 7.$$