

MA 137 – Calculus 1 with Life Science Applications
The Chain Rule and Higher Derivatives
(Sections 4.5 & 4.7)

Department of Mathematics
University of Kentucky

The Chain Rule

Theorem

If g is differentiable at x and f is differentiable at $y = g(x)$, then the composite function $(f \circ g)(x) = f[g(x)]$ is differentiable at x , and the derivative is given by

$$(f \circ g)'(x) = f'[g(x)] \cdot g'(x)$$

- The proof of the theorem is on p. 164 of the Neuhauser's textbook.
- The function g is the inner function; the function f is the outer function.
- The expression $f'[g(x)] \cdot g'(x)$ thus means that we need to find the derivative of the outer function, evaluated at $g(x)$, and the derivative of the inner function, evaluated at x , and then multiply the two together.
- A special case of the chain rule is called the **power chain rule**:

$$\text{If } y = [f(x)]^n \quad \text{then} \quad \frac{dy}{dx} = n[f(x)]^{n-1} \cdot f'(x)$$

The Chain Rule in Leibniz Notation

The derivative of $f \circ g$ can be written in Leibniz notation.

If we set $u = g(x)$, then

$$\begin{aligned} \frac{d}{dx}[(f \circ g)(x)] &= \frac{d}{dx}f[g(x)] \\ &\stackrel{u=g(x)}{=} \frac{d}{dx}f(u) \\ &= \frac{df}{du} \cdot \frac{du}{dx} \end{aligned}$$

This form of the chain rule emphasizes that, in order to differentiate $f \circ g$, we multiply the derivative of the outer function and the derivative of the inner function, the former evaluated at u , the latter at x .

Example 1: (Online Homework HW13, # 3)

Let $F(x) = f(f(x))$ and $G(x) = (F(x))^2$ and suppose that

$$f(5) = 3 \quad f(7) = 5 \quad f'(5) = 8 \quad f'(7) = 13$$

Find $F'(7)$ and $G'(7)$.

Example 2: (Online Homework HW13, # 6)

Let $f(x) = \frac{9}{(2x^2 - 3x + 6)^4}$. Find $f'(x)$.

The Quotient Rule Using the Chain Rule

We can prove quotient rule using the product and (power) chain rules. Treat the quotient f/g as a product of f and the reciprocal of g . I.e.,

$$\frac{f(x)}{g(x)} = f(x) \cdot g(x)^{-1}.$$

Next, apply the product rule

$$\left(\frac{f(x)}{g(x)}\right)' = [f(x) \cdot g(x)^{-1}]' = f'(x) \cdot g(x)^{-1} + f(x) \cdot [g(x)^{-1}]'$$

and apply the (power) chain rule to find $[g(x)^{-1}]'$. We obtain

$$= f'(x) \cdot [g(x)^{-1}] + f(x) \cdot [(-1)g(x)^{-2} \cdot g'(x)].$$

Finish by writing the expression with a common denominator of $[g(x)]^2$

$$= \frac{f'(x) \cdot g(x) - f(x) \cdot g'(x)}{[g(x)]^2}.$$

Example 3: (Neuhauser, Example # 5, p. 170)

Find the derivative of $h(x) = \left(\frac{x}{x+1}\right)^2$.

Example 4: (Neuhauser, Problem # 32, p. 173)

Differentiate $g(N) = \frac{N}{(k + bN)^3}$ with respect to N .

Assume that b and k are positive constants.

Example 5: (Neuhauser, Problem # 39, p. 174)

Find the derivative of

$$\frac{[f(x)]^2}{g(2x) + 2x}$$

assuming that f and g are both differentiable functions.

Higher Derivatives

- The derivative of a function f is itself a function. We refer to this derivative as the **first derivative**, denoted f' . If the first derivative exists, we say that the function is once differentiable.
- Given that the first derivative is a function, we can define its derivative (where it exists). This derivative is called the **second derivative** and is denoted f'' . If the second derivative exists, we say that the original function is twice differentiable.
- This second derivative is again a function; hence, we can define its derivative (where it exists). The result is the **third derivative**, denoted f''' . If the third derivative exists, we say that the original function is three times differentiable.
- We can continue in this manner; **from the fourth derivative on**, we denote the derivatives by $f^{(4)}$, $f^{(5)}$, and so on. If the n th derivative exists, we say that the original function is n times differentiable.

- Polynomials are functions that can be differentiated as many times as desired. The reason is that the first derivative of a polynomial of degree n is a polynomial of degree $n - 1$. Since the derivative is a polynomial as well, we can find its derivative, and so on. Eventually, the derivative will be equal to 0.
- We can write higher-order derivatives in Leibniz notation: The n th derivative of $f(x)$ is denoted by

$$\frac{d^n f}{dx^n}$$

Example 6: (Online Homework HW13, # 4)

Find the first and second derivatives of the following function

$$f(x) = (5 - 3x^2)^4$$

Example 7: (Online Homework HW13, # 16)

Find the first and second derivatives of the following function

$$y = \frac{1 - 4u}{1 + 3u}$$

Velocity and Acceleration

The velocity of an object that moves on a straight line is the derivative of the object's position. The derivative of the velocity is the acceleration.

If $s(t)$ denotes the position of an object moving on a straight line, $v(t)$ its velocity, and $a(t)$ its acceleration, then the three quantities are related as follows:

$$v(t) = \frac{ds}{dt} \quad \text{and} \quad a(t) = \frac{dv}{dt} = \frac{d^2s}{dt^2}.$$

Example 8: (Neuhauser, Problem # 15, p. 184)

Neglecting air resistance, the height h (in meters) of an object thrown vertically from the ground with initial velocity v_0 is given by

$$h(t) = v_0 t - \frac{1}{2} g t^2$$

where $g = 9.81\text{m/s}^2$ is the earth's gravitational constant and t is the time (in seconds) elapsed since the object was released.

- (a) Find the velocity and the acceleration of the object.
- (b) Find the time when the velocity is equal to 0. In which direction is the object traveling right before this time? in which direction right after this time?