

# MA 137 — Calculus 1 with Life Science Applications

## Antiderivatives

### (Section 5.10)

Department of Mathematics  
University of Kentucky

## From Differential to Integral Calculus

Roughly speaking, Calculus has two parts:

differential calculus and integral calculus

At the core of **differential calculus** (which we have been studying so far) is the concept of the instantaneous rate of change of a function. We have seen how this concept can be used to locally approximate functions, to identify maxima and minima, to decide stability of equilibria, etc.

**Integral calculus**, on the other hand, deals with accumulated change, and, thereby, recovering a function from a mathematical description of its instantaneous rate of change. This recovery process, interestingly enough, is related to the concept of finding the area enclosed by a curve. This will be studied in Chapter 6 (and in the follow up course, MA 138).

## Antiderivatives

Many mathematical operations have an inverse. For example, to undo addition we use subtraction. To undo exponentiation we take logarithms. The process of differentiation can be undone by a process called *antidifferentiation*.

To motivate antidifferentiation, suppose we know the rate at which a bacteria population is growing and want to know the size of the population at some future time. The problem is to find a function  $F$  whose derivative is a known function  $f$ .

### Definition

A function  $F$  is called an **antiderivative** of  $f$  on an interval  $I$  if  $F'(x) = f(x)$  for all  $x \in I$ .

**Warning:** Although we will learn rules that allow us to compute antiderivatives, this process is typically **much more** difficult than finding derivatives; in addition, there are even cases where it is impossible to find an expression for an antiderivative.

## Corollaries of MVT

Two corollaries of the Mean Value Theorem will help us in finding antiderivatives. The first one is Corollary 2 from Section 5.1 (p. 222 of Neuhäuser's textbook):

### Corollary 2

If  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , with  $f'(x) = 0$  for all  $x \in (a, b)$ , then  $f$  is constant on  $[a, b]$ .

Corollary 2 is the converse of the fact that  $f'(x) = 0$  whenever  $f(x)$  is a constant function. Corollary 2 tells us that all antiderivatives of a function that is identically 0 are constant functions.

Corollary 3 says that functions with identical derivative differ only by a constant; that is, to find all antiderivatives of a given function, we need only find one.

### Corollary 3

If  $F(x)$  and  $G(x)$  are antiderivatives of the continuous function  $f(x)$  on an interval  $I$ , then there exists a constant  $c$  such that  $G(x) = F(x) + c$  for all  $x \in I$ .

**Proof:** Since  $F(x)$  and  $G(x)$  are both antiderivatives of  $f(x)$ , it follows that  $F'(x) = f(x) = G'(x)$  for all  $x \in I$ . Thus

$$[F(x) - G(x)]' = F'(x) - G'(x) = f(x) - f(x) = 0.$$

It follows from Corollary 2, applied to the function  $F - G$ , that  $F(x) - G(x) = c$ , where  $c$  is a constant.



## The Indefinite Integral

### Notation

The *indefinite integral* of  $f(x)$ , denoted by

$$\int f(x) dx$$

represents the *general* antiderivative of  $f(x)$ .

For example,  $\int 3x^2 dx = x^3 + c$ , where  $c$  is any constant.

### Rules for Indefinite Integrals

A.  $\int k f(x) dx = k \int f(x) dx$   $k$  any constant

B.  $\int [f(x) \pm g(x)] dx = \left[ \int f(x) dx \right] \pm \left[ \int g(x) dx \right]$

5/15

## Basic Indefinite Integrals

The formulas below can be verified by differentiating the righthand side of each expression. The quantities  $a$  and  $c$  below denote (nonzero) constants.

1.  $\int x^n dx = \frac{1}{n+1} x^{n+1} + c$   $n \neq -1$

2.  $\int \frac{1}{x} dx = \ln |x| + c$

3.  $\int e^{ax} dx = \frac{1}{a} e^{ax} + c$

4.  $\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$

5.  $\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$

**Warning:** We do not have simple derivative rules for products and quotients, so we should not expect simple integral rules for products and quotients.

6/15

### Example 1: (Online Homework HW22, # 2)

Find the antiderivative  $F$  of  $f(x) = 5x^4 - 2x^5$  that satisfies  $F(0) = -10$ .

7/15

In other words:

$$\begin{aligned} F(x) &= \int (5x^4 - 2x^5) dx = \\ &= 5 \int x^4 dx - 2 \int x^5 dx \\ &= 5 \cdot \left[ \frac{1}{5} x^5 \right] - 2 \left[ \frac{1}{6} x^6 \right] + C \\ &= \underline{x^5 - \frac{1}{3} x^6 + C} \end{aligned}$$

describes all antiderivatives of  $f(x)$ .

We want the one such that  $F(0) = -10$ . Thus

$$-10 = F(0) = 0^5 - \frac{1}{3} 0^6 + C \quad \therefore \boxed{C = -10}$$

$$\therefore \boxed{F(x) = x^5 - \frac{1}{3} x^6 - 10}$$



**Example 2:**

Evaluate the indefinite integral  $\int (t^3 + 3t^2 + 4t + 9) dt$ .

$$\int (t^3 + 3t^2 + 4t + 9) dt$$

$$= \int t^3 dt + 3 \int t^2 dt + 4 \int t dt + 9 \int 1 \cdot dt$$

$$= \frac{1}{4} t^4 + 3 \cdot \left(\frac{1}{3} t^3\right) + 4 \left(\frac{1}{2} t^2\right) + 9 \cdot t + C$$

$$= \frac{1}{4} t^4 + t^3 + 2t^2 + 9t + C$$


---

8/15

**Example 3:** (Online Homework HW22, # 5)

Evaluate the indefinite integral  $\int x(10 - x^4) dx$ .

$$\int \underbrace{x \cdot (10 - x^4)} dx$$

there are no rules for the antiderivative of a product

$$= \int (10x - x^5) dx = 10 \int x dx - \int x^5 dx$$

$$= 10 \left(\frac{1}{2} x^2\right) - \left(\frac{1}{6} x^6\right) + C$$

$$= 5x^2 - \frac{1}{6} x^6 + C$$

9/15

**Example 4:** (Online Homework HW22, # 7)

Evaluate the indefinite integral  $\int \frac{9u^4 + 7\sqrt{u}}{u^2} du$ .

$$\int \frac{9u^4 + 7\sqrt{u}}{u^2} du$$

there are no rules for the antiderivative of a quotient ...


$$\begin{aligned} &= \int \left( \frac{9u^4}{u^2} + \frac{7\sqrt{u}}{u^2} \right) du = \int (9u^2 + 7u^{\frac{1}{2}-2}) du \\ &= \int (9u^2 + 7u^{-3/2}) du = 9 \int u^2 du + 7 \int u^{-3/2} du \\ &= 9 \left( \frac{1}{3} u^3 \right) + 7 \left( \frac{1}{-3/2+1} u^{-3/2+1} \right) + C = \\ &= 3u^3 + 7 \left( \frac{1}{-1/2} u^{-1/2} \right) + C = 3u^3 - 14 \frac{1}{\sqrt{u}} + C \\ &= \boxed{\frac{3u^3\sqrt{u} - 14}{\sqrt{u}} + C} \end{aligned}$$

**Example 5:** (Online Homework HW22, # 10)

Find a function  $f$  such that  $f'(x) = 4x^3$  and the line  $x + y = 0$  is tangent to the graph of  $f$ .

$$f'(x) = 4x^3 \implies f(x) = x^4 + C$$

We also know that "somewhere" this function is tangent to the line  $y = -x$ .



all vertical translates of  $y = x^4$

I.e. at some point of  $f(x) = x^4 + C$  the tangent line has slope  $-1$ .

So  $f'(x) = 4x^3 = -1 \iff x = -\sqrt[3]{\frac{1}{4}} \approx -0.62996$   
 what is the value of  $f$  at  $x = -0.62996$ ?  
 It must be the value of the tangent line at



that point!

$$\text{Hence } f(-0.62996) = 0.62996$$

since the tangent line is  $y = -x$ .

Hence  $f(x) = x^4 + C$  is such that

$$0.62996 = (-0.62996)^4 + C$$

$$\therefore C = 0.47247$$

Hence: 
$$f(x) = x^4 + 0.47247$$

**Example 6:** (Online Homework HW22, # 13)

Find  $f$  if  $f'''(x) = \sin(x)$ ,  $f(0) = 8$ ,  $f'(0) = 4$ , and  $f''(0) = -10$ .

$$f'''(x) = \sin x, \quad f(0) = 8, \quad f'(0) = 4, \quad f''(0) = -10$$

Now:  $f'''(x) = \sin x \implies f''(x) = -\cos x + C_1$

Since  $f''(0) = -10$  we have  $-10 = \underbrace{-\cos(0)}_1 + C_1$

$$\therefore C_1 = -10 + 1 = -9$$

So  $f''(x) = -\cos x - 9$ . Thus

$$f'(x) = -\sin x - 9x + C_2$$

Since  $f'(0) = 4$  we have  $4 = f'(0) = \underbrace{-\sin(0)}_0 + C_2$

$$\therefore C_2 = 4. \quad \text{Hence}$$

$$f'(x) = -\sin x - 9x + 4$$

Since  $f'(x) = -\sin x - 9x + 4$

we have that

$$f(x) = \cos x - \frac{9}{2}x^2 + 4x + C_3$$

Since  $f(0) = 8$  we have

$$8 = f(0) = \underbrace{\cos(0)}_1 + C_3$$

$$\therefore C_3 = 8 - 1 = 7$$

Finally

$$f(x) = \cos x - \frac{9}{2}x^2 + 4x + 7$$



## Solving Simple Differential Equations

In this course, we have repeatedly encountered differential equations ( $\equiv$  DEs). Occasionally, we showed that a certain function would solve a given differential equation.

What we learned so far translates into solving DEs of the form

$$\frac{dy}{dx} = f(x).$$

That is, the rate of change of  $y$  with respect to  $x$  depends only on  $x$ . We now know that if we can find one such function  $y$  such that  $y' = f(x)$ , then there is a whole family of functions with this property, all related by vertical translations.

If we want to pick out one of these functions, we need to specify an initial condition — a point  $(x_0, y_0)$  on the graph of the function. Such a function is called a solution of the **initial-value problem**

$$\frac{dy}{dx} = f(x) \quad \text{with } y = y_0 \text{ when } x = x_0.$$

13/15

## Example 7: (Neuhauser, Example 5, p. 298)

Solve the initial-value problem  $\frac{dy}{dx} = -2x^2 + 3$  with  $y_0 = 10$  when  $x_0 = 3$ .

14/15

$$\frac{dy}{dx} = -2x^2 + 3 \iff y = -\frac{2}{3}x^3 + 3x + C$$

Now when  $x_0 = 3$   $y_0 = 10$ . So

$$10 = -\frac{2}{3}(3)^3 + 3(3) + C$$

$$\iff$$

$$10 = -2 \cdot 9 + 9 + C$$

$$10 = -9 + C$$

$$\therefore C = 19$$

$$\therefore \boxed{y = -\frac{2}{3}x^3 + 3x + 19} \quad (11)$$

Note that we could think of  
 $\frac{dy}{dx} = -2x^2 + 3$  as

$$dy = (-2x^2 + 3) dx$$

Now take the indefinite integral of both sides

$$\int 1 \cdot dy = \int (-2x^2 + 3) dx$$

$$y + C_1 = -\frac{2}{3}x^3 + 3x + C_2$$

hence  
 $y = -\frac{2}{3}x^3 + 3x + \underbrace{(C_2 - C_1)}_{C \text{ constant}}$



**Example 8:**

What about finding the solution of the initial-value problem

$$\frac{dy}{dx} = ry \quad \text{with } y(0) = y_0 \text{ and } r \text{ a constant? How can we do it?}$$

15/15

We have already solved this differential equation while studying Section 5.1.

Now we give a nicer proof:

$$\frac{dy}{dx} = ry$$



$$\frac{1}{y} \cdot \frac{dy}{dx} = r$$

What is this? By the chain rule

$$\frac{d}{dx}(\ln y) = r$$

If the derivative of a function is a constant

then the antiderivative is

$$\frac{d}{dx}[\ln y] = r \iff \ln y = rx + C$$

Hence, by taking the exponential of both sides,

$$e^{\ln y} = e^{rx + C} \iff y = e^{rx} \cdot e^C$$

When  $x=0$  we have  $y(0) = y_0$  so

$$y_0 = y(0) = \underbrace{e^{r \cdot 0}}_1 \cdot e^C \quad \therefore e^C = y_0$$

Hence the solution is:

$$y(x) = y_0 e^{rx}$$

As we did in Example 6, we could think

$$\text{of } \frac{dy}{dx} = ry \quad \text{as}$$

$$\frac{1}{y} dy = r dx$$

That is we separated the variables; now we take the indefinite integral of both sides

$$\int \frac{1}{y} dy = \int r dx$$

$$\text{hence } \boxed{\ln y = rx + C}$$

now proceed as before....