

FastTrack — MA 137/MA 113 — BioCalculus

Functions (3): The Algebra of Functions

Alberto Corso – (alberto.corso@uky.edu)

Department of Mathematics – University of Kentucky

Goal: We learn how two functions can be combined to form new functions. We then define one-to-one functions, which allows us to introduce the notion of inverse of a one-to-one function. These topics are of importance when we study exponential and logarithmic functions.

1/29

Note

Consider the above definition $(f+g)(x) = f(x)+g(x)$.

The $+$ on the left hand side stands for the operation of addition of functions.

The $+$ on the right hand side, however, stands for addition of the numbers $f(x)$ and $g(x)$.

Similar remarks hold true for the other definitions.

3/29

Combining functions

Let f and g be functions with domains A and B . We define new functions $f + g$, $f - g$, fg , and f/g as follows:

$$(f + g)(x) = f(x) + g(x) \quad \text{Domain } A \cap B$$

$$(f - g)(x) = f(x) - g(x) \quad \text{Domain } A \cap B$$

$$(fg)(x) = f(x)g(x) \quad \text{Domain } A \cap B$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} \quad \text{Domain } \{x \in A \cap B \mid g(x) \neq 0\}$$

2/29

Example 1:

Let us consider the functions $f(x) = x^2 - 2x$ and $g(x) = 3x - 1$.

Find $f + g$, $f - g$, fg , and f/g and their domains.

4/29

Example 2:


Let us consider the functions $f(x) = \sqrt{9 - x^2}$ and $g(x) = \sqrt{x^2 - 1}$.

Find $f + g$, $f - g$, fg , and f/g and their domains.

5/29

Example 3:

Use graphical addition to sketch the graph of $f + g$.

graph of $f + g$

7/29

The graph of the function $f + g$ can be obtained from the graphs of f and g by **graphical addition**.

This means that to obtain the value of $f + g$ at any point x we add the corresponding values of $f(x)$ and $g(x)$, that is, the corresponding y -coordinates.

Similar statements can be made for the other operations on functions.

6/29

Composition of Functions



Given any two functions f and g , we start with a number x in the domain of g and find its image $g(x)$. If this number $g(x)$ is in the domain of f , we can then calculate the value of $f(g(x))$.

The result is a new function $h(x) = f(g(x))$ obtained by substituting g into f . It is called the *composition* (or *composite*) of f and g and is denoted by $f \circ g$ (read: 'f composed with g' or 'f after g')

$$(f \circ g)(x) \stackrel{\text{def}}{=} f(g(x)).$$


WARNING: $f \circ g \neq g \circ f$.

8/29



Example 5:

Let f and g be the functions considered in Example 3. Use the information provided by the graphs of f and g to find $f(g(1))$, $g(f(0))$, $f(g(0))$, and $g(f(4))$.



Example 4:

Use $f(x) = 3x - 5$ and $g(x) = 2 - x^2$ to evaluate:

$$f(g(0)) = \qquad g(f(0)) =$$

$$f(f(4)) = \qquad (g \circ g)(2) =$$

$$(f \circ g)(x) = \qquad (g \circ f)(x) =$$

Example 6:

Let $f(x) = \frac{x}{x+1}$ and $g(x) = 2x - 1$.

Find the functions $f \circ g$, $g \circ f$, and $f \circ f$ and their domains.


Example 7:

Express the function $F(x) = \frac{x^2}{x^2 + 4}$ in the form $F(x) = f(g(x))$.

Definition of a One-One Function

A function f with domain A is called a **one-to-one function** if no two elements of A have the same image, that is,
 $f(x_1) \neq f(x_2)$ whenever $x_1 \neq x_2$.

An equivalent way of writing the above condition is:
If $f(x_1) = f(x_2)$, then $x_1 = x_2$.



Example 8:

Find functions f and g so that $f \circ g = H$ if $H(x) = \sqrt[3]{2 + \sqrt{x}}$.

Horizontal Line Test



For functions that can be graphed in the coordinate plane, there is a useful criterion to determine whether a function is one-to-one or not.

Horizontal Line Test

A function is one-to-one



no horizontal line intersects its graph more than once.



Example 9:

Show that the function $f(x) = 5 - 2x$ is one-to-one.

17/29

<http://www.ms.uky.edu/~ma137>

Lecture #3

The Inverse of a Function


One-to-one functions are precisely those for which one can define a (unique) **inverse function** according to the following definition.

Definition of the Inverse of a Function

Let f be a one-to-one function with domain A and range B . Its **inverse function** f^{-1} has domain B and range A and is defined by

$$f^{-1}(y) = x \iff f(x) = y,$$

for any $y \in B$.



If f takes x to y ,
 then f^{-1} takes y back to x .
 I.e., f^{-1} undoes what f does.

NOTE:
 f^{-1} does NOT mean $\frac{1}{f}$.

19/29

<http://www.ms.uky.edu/~ma137>

Lecture #3

Example 10:

Graph the function $f(x) = (x - 2)^2 - 3$. The function is not one-to-one: Why? Can you restrict its domain so that the resulting function is one-to-one? (There is more than one correct answer.)

18/29

<http://www.ms.uky.edu/~ma137>

Lecture #3

Example 11:

Suppose $f(x)$ is a one-to-one function.

If $f(2) = 7$, $f(3) = -1$, $f(5) = 18$, $f^{-1}(2) = 6$ find:

$$f^{-1}(7) = \qquad f(6) =$$

$$f^{-1}(-1) = \qquad f(f^{-1}(18)) =$$

If $g(x) = 9 - 3x$, then $g^{-1}(3) =$

20/29

<http://www.ms.uky.edu/~ma137>

Lecture #3

Properties of Inverse Functions

Let $f(x)$ be a one-to-one function with domain A and range B . The inverse function $f^{-1}(x)$ satisfies the following "cancellation" properties:

$$f^{-1}(f(x)) = x \quad \text{for every } x \in A$$

$$f(f^{-1}(x)) = x \quad \text{for every } x \in B$$

Conversely, any function $f^{-1}(x)$ satisfying the above conditions is the inverse of $f(x)$.

Example 13:

Show that the functions $f(x) = \frac{1+3x}{5-2x}$ and $g(x) = \frac{5x-1}{2x+3}$ are inverses of each other.

Example 12:

Show that the functions $f(x) = x^5$ and $g(x) = x^{1/5}$ are inverses of each other.

How to find the Inverse of a One-to-One Function

1. Write $y = f(x)$.
2. Solve this equation for x in terms of y (if possible).
3. Interchange x and y . The resulting equation is $y = f^{-1}(x)$.

Example 14:

Find the inverse of $y = 4x - 7$.

Example 16:

Find the inverse of $y = \frac{2-x}{x+2}$.

Example 15:

Find the inverse of $y = \frac{1}{x+2}$.

Graph of the Inverse Function


The principle of interchanging x and y to find the inverse function also gives us a method for obtaining the graph of f^{-1} from the graph of f . **The graph of f^{-1} is obtained by reflecting the graph of f in the line $y = x$.**

The picture on the right hand side shows the graphs of:

$$f(x) = \sqrt{x+4}$$

and

$$f^{-1}(x) = x^2 - 4, \quad x \geq 0.$$



Example 17:

Find the inverse of the function $f(x) = 1 + \sqrt{1+x}$.
Find the domain and range of f and f^{-1} . Graph f and f^{-1} on the same cartesian plane.