

MA 138 – Calculus 2 with Life Science Applications  
**Functions of Two or More Independent Variables**  
(Section 10.1)

**Alberto Corso**

⟨alberto.corso@uky.edu⟩

Department of Mathematics  
University of Kentucky

## Coordinate Systems (in $\mathbb{R}^2$ and $\mathbb{R}^3$ )

Any point  $P$  in the plane can be represented as an ordered pair of real numbers. To locate a point in space, three numbers are required.


We first choose a fixed point  $O$  (the origin) and three directed lines through  $O$  that are perpendicular to each other, called the coordinate axes and labeled the  $x$ -axis,  $y$ -axis, and  $z$ -axis. Usually we think of the  $x$ - and  $y$ -axes as being horizontal and the  $z$ -axis as being vertical. The direction of the  $z$ -axis is determined by the right-hand rule: If you curl the fingers of your right hand around the  $z$ -axis in the direction of a  $90^\circ$  counterclockwise rotation from the positive  $x$ -axis to the positive  $y$ -axis, then your thumb points in the positive direction of the  $z$ -axis.

The three coordinate axes determine three coordinate planes: The  $xy$ -plane is the plane that contains the  $x$ - and  $y$ -axes; the  $yz$ -plane contains the  $y$ - and  $z$ -axes; the  $xz$ -plane contains the  $x$ - and  $z$ -axes.

These three coordinate planes divide space into eight parts, called octants. The first octant is determined by the positive axes.

Now if  $P$  is any point in space, let  $x_P$  be the distance from  $P$  to the  $yz$ -plane, let  $y_P$  be the distance from  $P$  to the  $xz$ -plane, and let  $z_P$  be the distance from  $P$  to the  $xy$ -plane. We represent the point  $P$  by the ordered triple  $(x_P, y_P, z_P)$  of real numbers and we call them the coordinates of  $P$ .

The point  $P(x_P, y_P, z_P)$  determines a rectangular box. Dropping a straight line from  $P$  to the  $xy$ -plane, we obtain the projection of  $P$  onto the  $xy$ -plane:  $Q(x_P, y_P, 0)$ . Similarly,  $R(0, y_P, z_P)$  and  $S(x_P, 0, z_P)$  are the projections of  $P$  onto the  $yz$ -plane and  $xz$ -plane, respectively.



## Example 1 (Problems # 7, 8, Section 10.1, p. 572)

- Locate the following points in a three-dimensional Cartesian coordinate system:

$$A(1, 3, 2) \quad B(-1, -2, 1) \quad C(0, 1, 2) \quad D(2, 0, 3)$$

- Describe the set of all points in  $\mathbb{R}^3$  that satisfy the following expressions:

$$(a) x = 0 \quad (b) y = 0 \quad (c) z = 0 \quad (d) z \geq 0 \quad (e) y \leq 0$$

# Functions of Two or More Independent Variables

We consider functions for which

- the **domain** consists of pairs of real numbers  $(x, y)$  with  $x, y \in \mathbb{R}$  or, more generally, of  $n$ -tuples of real numbers  $(x_1, x_2, \dots, x_n)$  with  $x_1, x_2, \dots, x_n \in \mathbb{R}$ . We write  $\mathbb{R}^n$  to denote the set of all  $n$ -tuples of real numbers  $(x_1, x_2, \dots, x_n)$ .
- the **range** consists of subsets of the real numbers.

## Real-Valued Functions

Suppose  $D \subset \mathbb{R}^n$ . Then a real-valued function  $f$  on  $D$  assigns a real number to each element in  $D$ , and we write

$$f: D \longrightarrow \mathbb{R}, \quad (x_1, x_2, \dots, x_n) \mapsto f(x_1, x_2, \dots, x_n)$$

The set  $D$  is the **domain** of the function  $f$ , and the set

$$\{w \in \mathbb{R} \mid f(x_1, x_2, \dots, x_n) = w \text{ for some } (x_1, x_2, \dots, x_n) \in D\}$$

is the **range** of the function  $f$ .

## Graph of a Function of Two Variables


- If  $f$  is a function of two independent variables, we usually denote the independent variables by  $x$  and  $y$ , and write  $f(x, y)$ .
- We also write  $z = f(x, y)$  to make explicit the value taken on by  $f$  at the general point  $(x, y)$ . The variable  $z$  is the dependent variable.
- If a function  $f$  is given by a formula and no domain is specified, then the domain of  $f$  is understood to be the set of all pairs  $(x, y)$  for which the given expression is well-defined.
- To visualize a function of two variables we often consider its graph.

### Graph of a Function of Two Variables

The graph of a function  $f$  of two independent variables with domain  $D$  is the set of all points  $(x, y, z) \in \mathbb{R}^3$  such that  $z = f(x, y)$  for  $(x, y) \in D$ . That is, the graph of  $f$  is the set

$$\text{Graph}(f) = \{(x, y, z) \mid z = f(x, y) \text{ with } (x, y) \in D\}.$$

The graph of  $f(x, y)$  is therefore a surface in three-dimensional space, as illustrated, for example, by the following picture



which shows the graph of the function

$$f(x, y) = x e^{-x^2 - y^2}$$

over the square  $[-2, 2] \times [-2, 2]$ .

Graphing a surface in three-dimensional space is difficult. Fortunately, good computer software is now available that facilitates this task.

## Example 2 (Online Homework # 2)

Suppose  $f(x, y) = xy^2 + 7$ . Compute the following values

- $f(4, -2)$
- $f(-2, 4)$
- $f(t, 4t)$
- $\frac{f(x_0, y_0 + h) - f(x_0, y_0)}{h}$



### Example 3 (Online Homework # 3)

Find the domain of the following functions





- $f(x, y) = \ln(x + y)$
- $g(x, y) = \sqrt{x^2 y^3}$
- $h(x, y) = e^{-\frac{1}{x+y}}$
- $k(x, y) = x^2 + y^3$

## Example 4 (Online Homework # 4)

Match the equation of the surface

$$z = \sin x \quad x^2 + y^2 = 4 \quad xyz = 0 \quad x^2 + z^2 = 4$$

with one of the graphs below



## Level Curves (or Contour Lines)

Another way to visualize functions is with **level curves** or **contour lines**. This approach is used, for instance, in topographical maps.

There is a *subtle distinction* between level curves and contour lines, in that **level curves are drawn in the function domain** whereas **contour lines are drawn on the surface**.

This distinction is not always made, and often the two terms are used interchangeably. Our text almost exclusively uses level curves, for which we now give the precise definition:

### Level curves

Suppose that  $f: D \rightarrow R$ ,  $D \subset \mathbb{R}^2$ . Then the level curves of  $f$  comprise the set of points  $(x, y)$  in the  $xy$ -plane where the function  $f$  has a constant value; that is,  $f(x, y) = c$ .

## Graph of $z = e^{-x^2-y^2}$






FIGURE: topographical map



The picture on the **left** shows the mesh plot on the graph of the function; the picture on the **right** shows the contour lines on the graph.











The picture shows the level curves of the function  $z = e^{-x^2-y^2}$  in the  $xy$ -plane

## Example 5 (Online Homework # 5, 6, 7)

Match each of the following functions of two variables  $x$  and  $y$

$$f(x, y) = x^2 - 2 \quad g(x, y) = 3 - x + y \quad h(x, y) = |x| + |y| \quad k(x, y) = xye^{-x^2 - y^2}$$

with its graph (labeled A.-D.) and its level curves (labeled I.-IV.).



## Example 6 (Problem #4, Exam 3, Spring 2012)

Find the largest possible domain for  $f(x, y) = \ln(x - 2y^2)$ .

Determine explicitly the equations of the level curves  $f(x, y) = c$  and graph them in the domain of  $f$ .

## Example 7 (Problem # 47, Section 10.1, p. 574)

The picture below shows the oxygen concentration for Long Lake, Clear Water County (Minnesota). The water flea *Daphnia* can survive only if the oxygen concentration is higher than 3 mg/l. Suppose that you wanted to sample the *Daphnia* population in 1997 on days 180, 200, and 220. Below which depths can you be fairly sure not to find any *Daphnia*?

