

**MA 138 – Calculus 2 with Life Science Applications**  
**Linear Systems: Theory**  
(Section 11.1)

**Alberto Corso**

⟨alberto.corso@uky.edu⟩

Department of Mathematics  
University of Kentucky

## Systems of Differential Equations

- Suppose that we are given a set of variables  $x_1, x_2, \dots, x_n$ , each depending on an independent variable, say,  $t$ , so that

$$x_1 = x_1(t), \quad x_2 = x_2(t), \quad \dots, \quad x_n = x_n(t).$$

- Suppose also that the dynamics of the variables are linked by  $n$  differential equations ( $\equiv$ DEs) of the first-order; that is,

$$\begin{cases} \frac{dx_1}{dt} = g_1(t, x_1, x_2, \dots, x_n) \\ \vdots \\ \frac{dx_n}{dt} = g_n(t, x_1, x_2, \dots, x_n) \end{cases}$$

- This set of equations is called a **system of differential equations**.
- On the LHS are the derivatives of  $x_i(t)$  with respect to  $t$ . On the RHS is a function  $g_i$  that depends on the variables  $x_1, x_2, \dots, x_n$  and on  $t$ .

## Examples

### ■ Kermack & McKendrick epidemic disease model (SIR, 1927)

$$\begin{cases} \frac{dS}{dt} = -bSI \\ \frac{dI}{dt} = bSI - aI \\ \frac{dR}{dt} = aI \end{cases}$$

$S = S(t)$  = # of susceptible individuals

$I = I(t)$  = # of infected individuals

$R = R(t)$  = # of removed individuals ( $\equiv$  no longer susceptible)

$a, b$  = constant rates

### ■ Lotka-Volterra predator-prey model (1910/1920):

$$\begin{cases} \frac{dN}{dt} = rN - aPN \\ \frac{dP}{dt} = abPN - dP \end{cases}$$

$N = N(t)$  = prey density

$P = P(t)$  = predator density

$r$  = intrinsic rate of increase of the prey

$a$  = attack rate

$b$  = efficiency rate of predators in turning preys into new offsprings

$d$  = rate of decline of the predators

## Direction Field of a System of 2 Autonomous DEs

- Review the notion of the direction field of a DE of the first order  $dy/dx = f(x, y)$ . We encountered this notion just before Section 8.2 (Handout; Lectures 15 & 16).
- Consider, now a system of two autonomous differential equations

$$\begin{cases} \frac{dx}{dt} = g_1(x, y) \\ \frac{dy}{dt} = g_2(x, y) \end{cases}$$

- Assuming that  $y$  is also a function of  $x$  and using the chain rule, we can eliminate  $t$  and obtain the DE


$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{g_2(x, y)}{g_1(x, y)}$$

of which we can plot the direction field.

## Example (Lotka-Volterra)

Consider the system of DEs  $\frac{dx}{dt} = x - 4xy$  and  $\frac{dy}{dt} = 2xy - 3y$ .

The direction field of the differential equation  $\frac{dy}{dx} = \frac{(2x-3)y}{x(1-4y)}$  has been produced with the SAGE commands in Chapter 8.



Notice that the trajectories are closed curves. Furthermore, they all seem to revolve around the point  $P(3/2, 1/4)$ . This is the point where the factors  $2x - 3$  and  $1 - 4y$  of  $dy/dt$  and  $dx/dt$ , respectively, are both zero.

## Linear Systems of Differential Equations (11.1)

- We first look at the case when the  $g_i$ 's are linear functions in the variables  $x_1, x_2, \dots, x_n$  — that is,

$$\begin{cases} \frac{dx_1}{dt} = a_{11}(t)x_1 + \dots + a_{1n}(t)x_n + f_1(t) \\ \vdots \\ \frac{dx_n}{dt} = a_{n1}(t)x_1 + \dots + a_{nn}(t)x_n + f_n(t) \end{cases}$$

- We can write the linear system in matrix form as

$$\frac{d}{dt} \begin{bmatrix} x_1(t) \\ \vdots \\ x_n(t) \end{bmatrix} = \begin{bmatrix} a_{11}(t) & \dots & a_{1n}(t) \\ \vdots & & \vdots \\ a_{n1}(t) & \dots & a_{nn}(t) \end{bmatrix} \begin{bmatrix} x_1(t) \\ \vdots \\ x_n(t) \end{bmatrix} + \begin{bmatrix} f_1(t) \\ \vdots \\ f_n(t) \end{bmatrix}$$

and we call it an **inhomogeneous system of linear, first-order differential equations**.

- We can write our inhomogeneous system of linear, first-order differential equations as follows

$$\frac{dx}{dt} = A(t)\mathbf{x} + \mathbf{f}(t)$$

- We are mainly interested in the case when  $\mathbf{f}(t) = \mathbf{0}$ , that is,

$$\frac{dx}{dt} = A(t)\mathbf{x},$$

an **homogeneous** system of linear, first-order differential equations.


- Finally, we will study the case in which  $A(t)$  does not depend on  $t$

$$\frac{dx}{dt} = A\mathbf{x},$$

an **homogeneous system of linear, first-order differential equations with constant coefficients.**

## Example 1 (Problem #8, Exam 3, Spring 2013)

- (a) Verify that the functions  $x(t) = e^{4t} + 5e^{-t}$  and  $y(t) = 4e^{4t} - 5e^{-t}$  (whose graphs are given below) are solutions of the system of DEs



$$\begin{cases} \frac{dx}{dt} = & y \\ \frac{dy}{dt} = 4x + 3y \end{cases}$$

with  $x(0) = 6$  and  $y(0) = -1$ .

- (b) Rewrite the given system of DEs and its solutions in the form

$$\underbrace{\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}}_{\text{system of differential equations}}$$

$$\underbrace{\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix} e^{4t} + 5 \begin{bmatrix} \gamma \\ \delta \end{bmatrix} e^{-t}}_{\text{solutions}}$$

for appropriate choices of the constants  $a, b, c, d, \alpha, \beta, \gamma$ , and  $\delta$ .



## Specific Solutions of a Linear System of DEs

- Consider the system  $\frac{d\mathbf{x}}{dt} = A\mathbf{x}$ .

- We claim that the vector-valued function

$$\mathbf{x}(t) = \begin{bmatrix} v_1 e^{\lambda t} \\ v_2 e^{\lambda t} \end{bmatrix} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} e^{\lambda t}$$

where  $\lambda$ ,  $v_1$  and  $v_2$  are constants, is a solution of the given system of DEs, for an appropriate choice of values for  $\lambda$ ,  $v_1$ , and  $v_2$ .

- More precisely,  $\begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$  is an eigenvector of the matrix  $A$  corresponding to the eigenvalue  $\lambda$  of  $A$ .

# The Superposition Principle

## Principle

Suppose that

$$\begin{bmatrix} \frac{dx_1}{dt} \\ \frac{dx_2}{dt} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix}.$$

If  $\mathbf{y}(t) = \begin{bmatrix} y_1(t) \\ y_2(t) \end{bmatrix}$  and  $\mathbf{z}(t) = \begin{bmatrix} z_1(t) \\ z_2(t) \end{bmatrix}$

are solutions of the given system of DEs, THEN

$$\mathbf{x}(t) = c_1\mathbf{y}(t) + c_2\mathbf{z}(t)$$

is also a solution of the given system of DEs for any constants  $c_1$  and  $c_2$ .

# The General Solution

## Theorem

Let

$$\frac{dx}{dt} = Ax$$

where  $A$  is a  $2 \times 2$  matrix with **two real and distinct eigenvalues**  $\lambda_1$  and  $\lambda_2$  with corresponding eigenvectors  $\mathbf{v}_1$  and  $\mathbf{v}_2$ .

THEN

$$x(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t}$$

is the general solution of the given system of DEs.

The constants  $c_1$  and  $c_2$  depend on the initial condition.