

The goal of this project is to establish an amusing formula relating π to an infinite product, which is known as Wallis's product. (Of course this infinite product has to be defined as a limit.) The formula is:

$$\frac{\pi}{2} = \frac{2}{1} \cdot \frac{2}{3} \cdot \frac{4}{3} \cdot \frac{4}{5} \cdot \frac{6}{5} \cdot \frac{6}{7} \cdots \quad (1)$$

The right-hand side of this expression can also be written as

$$\lim_{n \rightarrow \infty} \prod_{k=1}^n \frac{(2k)^2}{(2k-1)(2k+1)}.$$

Though we have not used the symbol \prod before, you should be able to guess what it means, once I tell you that \prod is related to products in the same way that \sum is related to sums.

Your assignment is to carry out the steps below in order to see why (1) is true. The tools you will use are integration by parts (which is probably the most useful technique for studying integrals) and the comparison properties for integrals.

You should work in groups of 1 to 3 students. Each group will hand in one report for the group and all group members will receive the same grade. Each of you should participate in preparing the report and on the front of your paper please write "All group members participated in the preparation of this project". Each group member should sign this statement indicating they agree with this statement.

In your writeup, you should not just give the computations without explanation. The best papers will probably not give every detail of every step of the computation but will give a sentence or two describing what is being done and then the result of several steps. When you use results from the text (or from other sources) be sure to give a precise reference by page and equation number.

You should feel free to ask for help—tell us where you are stuck, and we will try to point you in the right direction. You are also free to consult with your fellow students.

However, you should not copy directly from another paper, obviously.

1. Derive the reduction formula for \cos , #34 inside the back cover of Varberg, Purcell and Rigdon. Hint: What I really wanted to do was ask you to derive the reduction formula for \sin , #73. Unfortunately, this is done for you in section 8.4. You should imitate the argument used in example 7 p. 389 to establish the reduction formula for \cos .

2. Use the reduction formula for \sin to show that for $n = 2, 3, \dots$,

$$\int_0^{\pi/2} \sin^n x \, dx = \frac{n-1}{n} \int_0^{\pi/2} \sin^{n-2} x \, dx.$$

3. Show that for $n = 1, 2, \dots$

$$\int_0^{\pi/2} \sin^{2n+1} x \, dx = \frac{2}{3} \cdot \frac{4}{5} \cdot \dots \cdot \frac{2n}{2n+1} = \prod_{k=1}^n \frac{2k}{2k+1}$$

and

$$\int_0^{\pi/2} \sin^{2n} x \, dx = \frac{\pi}{2} \cdot \frac{1}{2} \cdot \frac{3}{4} \cdot \dots \cdot \frac{2n-1}{2n} = \frac{\pi}{2} \prod_{k=1}^n \frac{2k-1}{2k}.$$

Hint: Before you try the general formula, you might want to try a few concrete examples, say $n = 0, 1$ and 2 . After you see the pattern, use mathematical induction to establish the formulae for all n .

4. Divide the two formulae in part 3 and show

$$\frac{\pi \int_0^{\pi/2} \sin^{2n+1} x \, dx}{2 \int_0^{\pi/2} \sin^{2n} x \, dx} = \frac{2}{1} \cdot \frac{2}{3} \cdot \frac{4}{3} \cdot \frac{4}{5} \cdot \frac{6}{5} \cdot \frac{6}{7} \cdot \dots \cdot \frac{2n}{2n-1} \cdot \frac{2n}{2n+1}.$$

5. To finish, we need to show that

$$\lim_{n \rightarrow \infty} \frac{\int_0^{\pi/2} \sin^{2n+1} x \, dx}{\int_0^{\pi/2} \sin^{2n} x \, dx} = 1. \quad (2)$$

To do this, we begin with the inequalities

$$\sin^{2n+2} x \leq \sin^{2n+1} x \leq \sin^{2n} x, \quad n = 0, 1, 2, \dots$$

For which x are these true? Why? Using these inequalities and part 2, show that

$$f(n) \leq \frac{\int_0^{\pi/2} \sin^{2n+1} x \, dx}{\int_0^{\pi/2} \sin^{2n} x \, dx} \leq h(n)$$

where

$$\lim_{n \rightarrow \infty} f(n) = \lim_{n \rightarrow \infty} h(n) = 1.$$

Now the squeeze theorem tells us that (2) and hence (1) holds. In your report, you should find explicit formulae for the functions f and h .

6. Who was Wallis and how did this product arise?

The above is adapted from the book *Calculus*, by Michael Spivak. In your copious spare time, you look through this book to see a different view of what a calculus book should be.